



## Marsupials

*Please note that these options are not recommendations for a particular taxon, but possible choices that depend on individual circumstances. It is the responsibility of the veterinarian and animal care staff to determine the dosage and best treatment for an individual based short-term and long-term reproductive goals, facility considerations, and logistical concerns.*

### THE CURRENT OPTIONS FOR FEMALES INCLUDE:

MGA implants (progestin)

Depo-Provera® injections (progestin)

Suprelorin® implants (GnRH agonist)

Lupron® injections (GnRH agonist)

Porcine zona pellucida (PZP) (immunocontraception)

*Note: GnRH agonists have been shown to be ineffective in male marsupials.*

*Note: PZP appears to be effective in marsupials but this information is based on limited data (n=6)*

### PROGESTINS

#### MGA Implants (melengestrol acetate)

- Duration of efficacy: Effective for at least 2 years, but may be much longer if not removed.
- Route of administration: Surgically inserted after proper sterilization (see [MGA Implant Product Page](#)).
- Latency to effectiveness: 1-2 weeks.
- Dosing: Doses vary by taxon. Dosing guidelines can be found on the [MGA Implant Product Page](#).
- Estrous cycles during contraceptive treatment: Unusual but can sometimes occur.
- Use during pregnancy: There is no evidence of problems during early pregnancy; however not recommended in late pregnancy because of the possibility of prolonged gestation, although the effect may depend on species and dose.
- Use during lactation: Can be used in lactating females.
- Use in seasonal breeders: Implants should be placed at least 1 month prior to breeding season.
- Reversibility: Generally reliable after implants are removed.
- Use in prepubertal animals: The lack of data on pre-pubertal treatment and potential long-term effects on fertility contraindicates recommending contraception before puberty.
- Behavioral effects: Data deficient.
- Effects on physical characteristics: Data deficient.
- Other: Please note that MGA implants are not available outside the U.S.

#### Depo-Provera® Injections (medroxyprogesterone acetate)

- Duration of efficacy: Extremely variable, with reports ranging from 1 to 24 months.
- Route of administration: Injectable.
- Latency to effectiveness: 1-2 weeks.
- Dosing: Doses and injection frequency vary by taxon. Dosing guidelines can be found on the [Depo-Provera Product Page](#).

- Estrous cycles during contraceptive treatment: Unusual but can sometimes occur.
- Use during pregnancy: Depo-Provera is not recommended for females that might be pregnant because its potentially long duration of efficacy may interfere with parturition; females should be confirmed not pregnant before starting treatment.
- Use during lactation: Can be used in lactating females.
- Use in seasonal breeders: First injection should be given at least 1 month prior to breeding season.
- Reversibility: Data from various mammal species have shown that duration of efficacy and latency to conception following last injection can be quite variable across individuals (from 1 month to 2 years); however, individuals do tend to reverse after treatment ends.
- Use in prepubertal animals: The lack of data on pre-pubertal treatment and potential long-term effects on fertility contraindicates recommending contraception before puberty.
- Behavioral effects: While there may be individual and species variation in response, females may show male-like qualities such as increased aggression.
- Effects on physical characteristics: While there may be individual and species variation in response, females may develop male-type secondary sex characteristics.

#### **Progestin-Related Cautions**

- Progestins may result in weight gain.
- In areas where cleaning is done by hosing (i.e., water under pressure), aerosolization of fecal matter, including excreted steroid hormones from MGA or Depo-Provera, is possible. Proper precautions should be taken.

#### **GNRH AGONISTS**

##### **Suprelorin® Implants (deslorelin acetate)**

- Duration of efficacy: 4.7mg implants are effective a minimum of 6 months, and 9.4mg implants are effective a minimum of 12 months but either formulation may be effective much longer; there is also individual and species variation in duration of efficacy.
- Route of administration: Subcutaneous via trocar in a place where it can be easily detected (e.g., base of the ear, inner thigh, or umbilical region) to facilitate removal when new ones are placed or when a breeding recommendation is received, even if implants are “expired”, to reduce duration of efficacy (see [Suprelorin Product Page](#)).
- Latency to effectiveness: About 3 weeks for females; due to the initial stimulation phase (see [Suprelorin Product Page](#)), implanted females must be separated from males for 3 weeks or oral megestrol acetate ([Ovaban or Megace](#)) must be used for 7 days prior through 7 days after implant placement to prevent the agonist-induced stimulation phase.
- Dosing: Doses vary by taxon. Dosage guidelines can be obtained by emailing the RMC at [contraception@stlzoo.org](mailto:contraception@stlzoo.org).
- Estrous cycles during contraceptive treatment: Estrus and ovulation are possible during the 3 weeks of stimulation, then down-regulation occurs. To prevent the stimulation phase, the oral megestrol acetate protocol described above is recommended. Some follicle growth may continue, even following down-regulation.
- Use during pregnancy: Not recommended; may cause abortion or if pregnancy progresses, mammary development may be suppressed impairing milk production.

- Use during lactation: No contraindications once lactation is established.
- Use in seasonal breeders: GnRH agonists can induce estrus in females even during the non-breeding season. When used in seasonal breeders, implants should be placed in females at least 1 month before the time of first seasonally anticipated estrus.
- Reversibility: Designed to be reversible, but time to reversal can be quite variable. To decrease time to reversal, implants should be removed.
- Behavioral effects: In general, the effects should be similar to those from ovariectomy or castration; possible increased appetite can result in weight gain unless food is restricted.
- Effects on physical characteristics: In general, the effects should be similar to those from ovariectomy.
- Other: The RMC is only able to distribute Suprelorin implants to AZA-accredited institutions or for animals managed under an SSP or Recovery SAFE Program. Suprelorin F® is commercially available in the U.S. through veterinarians, but solely for the treatment of ferret adrenal disease. For institutions outside of the U.S., Suprelorin is commercially available in the U.K., Europe, Australia, and New Zealand. To order implants, a [Suprelorin Registration Form](#), found on the [Suprelorin Product Page](#), must be submitted.
- **Caution:** Depo-Provera should not be substituted for oral megestrol acetate to suppress the Suprelorin stimulation phase because its prolonged initial high levels can interfere with Suprelorin-mediated down-regulation or the reproductive system.

#### **Lupron® Injections (leuprolide acetate)**

- **Caution:** Since data for Lupron are limited, most of this information is extrapolated from the use of Suprelorin.
- Duration of efficacy: Available in formulations for human use effective from 1 to 6 months, but duration of efficacy and time to reversal in wildlife species may vary.
- Route of administration: Injectable.
- Latency to effectiveness: About 3 weeks for females; due to the initial stimulation phase; implanted females must be separated from males for 3 weeks or oral megestrol acetate (Ovaban) must be used for 7 days prior through 7 days after implant placement to prevent the agonist-induced stimulation phase.
- Dosing: Doses vary by taxon. Dosage must be extrapolated from human dosing regimens due to limited use in other wildlife.
- Estrous cycles during contraceptive treatment: As with Suprelorin, estrus and ovulation may occur during the 3 weeks of stimulation, then down-regulation occurs. To prevent the stimulation phase, the oral megestrol acetate protocol described above is recommended.
- Use during pregnancy: Not recommended; may cause abortion or if pregnancy progresses, mammary development may be suppressed impairing milk production.
- Use during lactation: No contraindications once lactation is established.
- Use in seasonal breeders: GnRH agonists can induce estrus in females even during the non-breeding season. When used in seasonal breeders, implants should be placed in females at least 1 month before the time of first seasonally anticipated estrus.
- Reversibility: Duration of efficacy/time to reversal may vary.
- Use in prepubertal animals: The lack of data on pre-pubertal treatment and potential long-term effects on fertility contraindicates recommending contraception before puberty. Also, because

Lupron suppresses gonadal steroids, its use may delay epiphyseal closure of the long bones, resulting in taller individuals, similar to the effects of pre-pubertal spaying and neutering in domestic dogs and cats.

- Behavioral effects: In general, the effects should be similar to those from ovariectomy or castration; possible increased appetite can result in weight gain unless food is restricted.
- Effects on physical characteristics: In general, the effects should be similar to those from ovariectomy.
- Other: Lupron, approved for treatment of prostate cancer in men, is expensive but can sometimes be acquired through donation from the manufacturer. In wildlife, it has been used primarily in males to suppress testosterone and sperm production. It has rarely been used in zoo animals since the late 1990s due to cost.

### **GnRH-Agonist-Related Cautions**

- GnRH agonists have been shown to be ineffective in male marsupials.

## **IMMUNOCONTRACEPTION**

### **Porcine Zona Pellucida Vaccine**

- Duration of efficacy: Species-dependent; for most species, it is effective 6 months to 1 year. First and second injections should be given no sooner than 2 weeks apart and subsequent boosters administered as needed. For year-round breeders, boosters should be given every 8 months.
- Route of administration: Injectable intramuscular.
- Latency to effectiveness: Effective only after the primer and initial booster injection (typically given at 2-4 week intervals), depending upon species and adjuvant. There must be a minimum 2-week interval after the second dose before the male is placed with the female.
- Dosing: Doses vary by taxon. Dosage guidelines can be obtained from the Science and Conservation Center at <https://www.sccpzp.org> when ordering this product.
- Estrous cycles during contraceptive treatment: Animals will continue to show signs of breeding behavior even when they are adequately contracepted. The breeding season may extend beyond what is considered typical, resulting in additional estrous cycles. However, with repeated treatment, cycles may become irregular and eventually cease.
- Use during pregnancy: Does not interrupt pregnancy or affect fetal development.
- Use during lactation: No known contraindications based on data from feral horse mares; data deficient in other species.
- Use in prepubertal animals: The lack of data on pre-pubertal treatment contraindicates recommending contraception before puberty.
- Use in seasonal breeders: Because PZP is not effective until after at least 2 injections (typically given 2-4 weeks apart), depending on species and adjuvant, treatment should be initiated at least 2 months before the anticipated onset of the breeding season.
- Reversibility: Is intended to be reversible, but repeated treatment can extend time to reversal. Because reversal becomes less likely with repeated treatment, use should be limited to 3-4 consecutive years or to animals not essential for breeding programs.

- Behavioral effects: Since the vaccine usually doesn't suppress estrous cycles, it has little or no effect on social behavior. In some species the failure to conceive can result in longer than usual breeding season, and in some cases this can result in aggression and social disruption.
- Effects on physical characteristics: Data deficient.
- Other: Contact Kim Frank at the Science and Conservation Center ([kfrank@sccpzp.org](mailto:kfrank@sccpzp.org)) for more detailed instructions and to order this product.

#### **PZP-Related Cautions**

- PZP may not be reversible after long-term use (>3 consecutive years).

***For more details on each of these products, please refer the specific product page.***

***Reporting requirements:*** Any use of Suprelorin implants in the United States must be reported to the RMC via our online contraception survey website (<https://www.zoocontraceptiondata.org>), including any and all adverse events associated with product use. Additionally, in order to increase our knowledge of the efficacy and reversibility of other contraception products, it is recommended that all individuals on contraception be added to the RMC's contraception database via our online contraception survey website (<https://www.zoocontraceptiondata.org>).

#### ***Ongoing Studies for which sample collection is encouraged:***

- ***RHSP Archive*** - The RMC and the Reproductive Health Surveillance Program (RHSP) request that facilities submit complete reproductive tracts to the RHSP anytime an animal dies or has their reproductive tract removed, to be available for investigations of reproductive health. See the RHSP website ([www.stlzoo.org/RHSP](http://www.stlzoo.org/RHSP)) for more specifics.
- ***Deslorelin Assay Validation*** - The RMC requests that facilities using Suprelorin implants, which contain deslorelin as the active ingredient, collect serum samples any time the animal is in hand after implant placement to help us establish a database of effective deslorelin concentrations and dynamics.

***Disclaimer:*** The RMC strives to provide accurate and current contraception recommendations based on various sources (e.g., publications, AZA RMC/EAZA RMG database records); however, as these are prescription-only medicines, it is the responsibility of the veterinarian and animal managers to determine the dosage and best treatment for an individual.